

2

DTIC FILE COPY

AD-A200 388

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470-P00003

R&T Code NR 33359-718

Technical Report No. 109

Infrared Spectroelectrochemistry of Surface Species:
An in situ Surface Fourier Transform Infrared Study
of Adsorption of Isoquinoline at a Mercury Electrode

by

D. Blackwood, C. Korzeniewski, W. McKenna, J. Li and S. Pons

Prepared for publication in Mol. Phen. at Electrode Surf., ACS Symp. Series

Department of Chemistry
University of Utah
Salt Lake City, UT 84112

July 15, 1988

DTIC
ELECTE
NOV 14 1988
S D
E

Reproduction in whole, or in part, is permitted for
any purpose of the United States Government

This document has been approved
for publication and sale; its
distribution is unlimited.

28 11 10 08A

REPORT DOCUMENTATION PAGE

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release and sale. Distribution unlimited.		
2b DECLASSIFICATION / DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) ONR Technical Report No. 109			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION University of Utah		6b OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Henry Eyring Building Salt Lake City, UT 84112				7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-K-0470-P00003	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program, Code 1113 800 N. Quincy Street Arlington, VA 22217				10 SOURCE OF FUNDING NUMBERS	
				PROGRAM ELEMENT NO.	PROJECT NO.
11. TITLE (Include Security Classification) Infrared Spectroelectrochemistry of Surface Species: An in situ Surface Fourier Transform Infrared Study of Adsorption of Isoquinoline at a Mercury Electrode					
12. PERSONAL AUTHOR(S) D. Blackwood, C. Korzeniewski, W. McKenna, J. Li and S. Pons					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 9/87 TO 7/88		14. DATE OF REPORT (Year, Month, Day) July 15, 1988	
15. PAGE COUNT 24					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) infrared spectroelectrochemistry , mercury		
FIELD	GROUP	SUB-GROUP			
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Attached.					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a NAME OF RESPONSIBLE INDIVIDUAL Stanley Pons			22b TELEPHONE (Include Area Code) (801)581-4760		22c OFFICE

ABSTRACT

Infrared spectroscopy, and in particular the method of subtractively normalized interfacial Fourier transform infrared spectroscopy (SNIFTIRS), has been used extensively to examine interactions of species at the electrode/electrolyte interface. In most experiments to date, interactions at solid electrodes have been studied. A method to probe interactions at the mercury solution has been developed and is presented in this presentation. The potential dependent frequency shifts of species adsorbed at mercury electrodes are compared with shifts observed for similar species adsorbed on d-band metals.

Isosquintal (P. 11)

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	
Special	
A-1	



✓

INFRARED SPECTROELECTROCHEMISTRY OF SURFACE SPECIES:
AN IN SITU SURFACE FOURIER TRANSFORM INFRARED STUDY OF
ADSORPTION OF ISOQUINOLINE AT A STATIONARY MERCURY ELECTRODE.

Daniel Blackwood, Carol Korzeniewski¹, William McKenna,
Jianguo Li and Stanley Pons*

Department of Chemistry
University of Utah,
Salt Lake City UT 84112

¹Department of Chemistry
University of Michigan
Ann Arbor, MI

*To whom correspondence should be addressed.

INTRODUCTION

Even though the basic technique was developed over two decades ago, infrared spectroscopy of the electrode solution interface has only yielded really useful data in the last few years (1-5). During this time, three basic techniques have been developed. The first successful method was electrochemically modulated infrared spectroscopy, EMIRS (4), which is a dispersive technique in which the electrode potential is modulated at a set frequency (10 Hz) and the resulting attenuation of the reflecting infrared radiation is analyzed by phase sensitive detection. The second common infrared technique used in electrochemistry is infrared reflection absorption spectroscopy, IRRAS¹ (6), in which the polarization of the incident infrared radiation is rapidly modulated (74 kHz) between the s- and the p-states. As s-polarized light undergoes a 180° phase shift upon reflection from a metal surface, the electromagnetically induced standing wave develops a node at the surface so that there is effectively no field intensity. As a result, only p-polarized light can interact with species which are either adsorbed onto, or within a percentage of a wavelength of, the electrode surface. The perturbed infrared signal is again monitored with the aid of phase sensitive detection techniques. The third technique is subtractively normalized interfacial Fourier transform spectroscopy, SNIFTIRS (7). Here, pseudo steady state infrared spectra are collected at two different electrochemical states or potentials, so that the electrochemical difference spectra can be obtained simply by subtracting one from the other and normalizing.

In situ infrared reflection techniques can distinguish between absorption bands arising from solution species from those due to species adsorbed on the metal electrode surface by noting the rules listed in Table 1. These rules arise

due to the difference in the amount of phase shift undergone upon reflection by s- and p-polarized light as discussed above, and to the fact that the strength of the chemical bond between an adsorbed species and the electrode is expected to be a function of the electrode potential, as this will govern the amount of overlap that can occur between the bonding and antibonding orbitals of the adsorbate.

To illustrate the potential dependence of adsorbed species, the SNIFTIRS spectra of the b_{3u} ring bending mode of p-difluorobenzene at a platinum electrode is shown in Figure 1 (8). The upward pointing band is clearly independent of potential and was thus assigned to a solution species whereas the position of the downward pointing band shifts monotonically (at constant ionic strength) with potential and is therefore clearly due to some adsorbed species.

The particular system investigated for this presentation was the adsorption of isoquinoline on mercury surfaces, which has been previously studied by electrocapillary methods (9), ellipsometry (10), double layer capacity measurements (11), and a range of potential step techniques (12-15). The interest in this system is due in part to the fact that one observes well defined transitions in its physical properties as the adsorbed molecules undergo transitions in surface orientation and packing density under certain experimental conditions. Isoquinoline molecules has been shown to be adsorbed on mercury in four different orientations (Figure 2). The previous investigations indicate the following behavior for the isoquinoline orientation as a function of potential and concentration: at low negative potentials and low bulk concentrations, the molecules are believed to lie flat on the mercury's surface (molecular plane parallel to the surface). However, on increasing either the potential (in the negative direction) or the bulk concentration, the isoquinoline molecules are forced up into either the 4,5 position (10) or the 5,6 position (9). This

reorientation occurs gradually with the changing coordinates, and proceeds through a series of phases containing mixtures of these three orientations of isoquinoline molecules.

Increases in the potential to more negative values, and at sufficiently high concentrations, results in an abrupt reorientation to the 6,7 position. The reason that this second transition is much sharper than the first lies in the fact that mixed phases which would contain the 6,7 orientation are energetically less favorable than a complete monolayer of any of the pure standing orientational phases. Gierst et al (9) have produced a graph showing the dependence of the superficial excess on both potential and bulk concentration from their electrocapillary data; we reproduce some of their data in Figure 3.

For an isoquinoline molecule adsorbed onto the surface of mercury, the component of its total dipole moment that is perpendicular to the surface will increase as its orientation changes from:

Flat - 4,5 - 5,6 - 6,7

The surface selection rules for infrared radiation reflection from metal surfaces and its interactions with adsorbed species are such that infrared radiation should be absorbed most strongly by vibrational modes that have a component of the dipole derivative (with respect to the normal coordinate) normal to the metal surface. Therefore as the isoquinoline molecule reorients in the order listed above, the absorption of infrared radiation by the in-plane vibrational modes would be expected to increase, while that of the out-of-plane modes would be predicted to decrease. In the flat orientation there is no component of the dipole moment perpendicular to the surface for the in-plane modes, thus it may be thought that these modes will not be able to absorb any of the incident

radiation. However, infrared active modes (and in some cases infrared forbidden transitions) can still be observed due to field-induced infrared absorption (16-20).

EXPERIMENTAL

Isoquinoline (Aldrich 97%) was further purified by refluxing with BaO for 30 minutes and distilling under vacuum. The resulting white crystalline solid had a melting point of 26°C. The purified isoquinoline was stored in the dark, at 0°C and under an argon atmosphere. Mercury was triple distilled (American Scientific) and all other chemicals were of AnalaR grade quality. All solutions were prepared with triply distilled water. All glassware was cleaned in a 50:50 (v:v) mixture of HNO₃ and H₂SO₄, rinsed with triply distilled water and steamed (triply distilled water) for half an hour. Because water strongly absorbs infrared radiation a thin layer cell was designed (Figure 4) which could be mounted in a vertical position on the spectrometer. The mercury was held in position by a simple glass tube, and electrical contact was achieved with a piece of platinum wire, dipped directly into the mercury. All potentials reported are with reference to a saturated calomel electrode.

The technique used to acquire all the data shown in this paper was SNIFTIRS, a schematic diagram of the required apparatus is shown in Figure (5). The FTIR spectrometer used was a vacuum bench Bruker IBM Model IR/98, modified so that the optical beam was brought upwards through the sample compartment and made to reflect from the bottom of the horizontal mercury surface. The infrared study of mercury electrode surfaces has been impeded by experimental difficulties in cell design and optical considerations; the methods used herein are adapted from a configuration that has been used by Bewick and co-workers (21).

RESULTS AND DISCUSSION

The abrupt transition to the 6,7 orientation manifests itself in cyclic voltammetry as a sharp current spike (Figure 6). The cathodic spike was found to contain a charge of $2.9 \mu\text{C cm}^{-2}$, while its anodic counterpart contained $3.3 \mu\text{C cm}^{-2}$. The peak separation was 100 mV, although this relatively large value is due in part to the high iR drop present in thin layer cells.

Figure 7 shows typical SNIFTIRS spectra for isoquinoline molecules adsorbed on mercury. The reference spectrum in each case was obtained at 0.0V vs. SCE reference electrode; at this potential the molecules are believed to be oriented flat on the metal surface. The vibrational frequencies of the band structure (positive values of absorbance) are easily assigned since they are essentially the same as those reported by Wait et al.(22) for pure isoquinoline. The differences in the spectra are that the bands for the adsorbed species exhibit blue shifting of $3\text{-}4 \text{ cm}^{-1}$ relative to the neat material, and the relative intensities of the bands in each case are markedly changed.

The major vibrational modes observed for isoquinoline are listed in Table 2. The assignments made by Wait et al.(22) are also included. These authors made their assignments from considerations of the higher $D_{\infty h}$ symmetry parent species, instead of the C_s symmetry group; they demonstrated that the assignments arising from this representation are reasonable.

The difference spectra show a complete absence of bands with negative absorbances (Figure 7), this can be explained if the vibrational frequencies of the bands do not shift with changes in the electrode potential, and that they are adsorbed over the entire potential region investigated. This is consistent with results of electrochemical double layer experiments. The spectra can now be interpreted as relatively simple changes in the absorption of infrared radiation.

The intensities of the bands are markedly potential dependent; an especially large change in the intensities is observed at potentials where the orientation changes to the vertical 6,7 configuration.

In order to explain the lack of potential dependence of the vibrational frequencies of the bands it is necessary to examine how this characteristic of the vibrational modes of adsorbed species is believed to arise. Two types of mechanisms have been proposed to explain the potential dependent shift of vibrational frequencies. The first involves molecular orbital arguments; the second is based on arguments for interactions between the electric field across the double layer and the highly polarizable electrons of the adsorbed molecule (an electrochemical Stark effect). In the molecular orbital mechanism, electrons can be donated to empty metal orbitals through σ -type overlap with filled ligand orbitals of the appropriate symmetry. The metal can "back" donate electrons from filled d-orbitals to empty π^* antibonding orbitals on the adsorbate. When a molecule is adsorbed on a clean uncharged metal surface, its vibrational frequency may either increase or decrease from the frequency of the unadsorbed molecule depending upon the relative contributions of the σ - and π -bonding interactions. If the π -bonding interaction is dominant the frequency will decrease; conversely, the frequency will increase if the σ -bonding interaction is dominant. When the charge on the electrode is made negative, the bond is weakened due to donation of charge from the metal into adsorbate π^* orbitals and the band frequency shifts to lower wavenumber. When the charge on the metal is made positive a shift to higher frequency occurs. At a mercury electrode, however, there are no p- or d-electrons available to participate in a back-bonding interaction. The observation of potential dependent frequency shifts are therefore not expected according to this model.

The electric field mechanism involves coupling of the electric field across the double layer with highly polarizable electrons of the adsorbate. According to the Gouy-Chapman-Stern model, for high concentrations of supporting electrolytes, most of the potential drop will occur within the first 5-10Å of the electrode surface, and the drop will be approximately linear with distance. When a layer of adsorbed species is present, it can act as a dielectric across which the greatest portion of the potential drop will occur. Electric fields on the order of 10^9 V m^{-1} can exist in this region. Interaction of this electric field with the dipole moment of the molecule leads to changes in the vibrational frequency of the molecule.

The absorbances in Figure 7 have positive values. This indicates that the absorption of infrared radiation is strongest when the isoquinoline molecules are lying flat on the electrode surface (in these difference spectra, positive values of absorbance denote stronger absorption at the positive potential, i.e. potentials where the isoquinoline is adsorbed in the flat configuration). This is an opposite result than that suggested from the surface selection rule, and suggests that there is a strong field-induced absorption for the in-plane modes in this configuration, similar to that observed in previous work for pyrene adsorbed on platinum (18).

A closer examination of the SNIFTIRS difference spectra shows that there are marked differences in the changes in intensity of the in-plane and out-of-plane vibrational modes of the adsorbed isoquinoline with potential (Figure 8). The normalized intensities (against their intensity at -0.1V vs. SCE) of the vibrational bands shown in Figure 7 were plotted against potential (Figure 9). Since the bands in Figure 7 have positive values of absorbance, the positive vertical axis in Figure 9 represents a decrease in the amount of infrared

radiation absorbed. The figure clearly shows that the amount of radiation absorbed by the out-of-plane vibrational modes decreases up to a factor of 10 as the potential is changed from -0.1 to -1.0V, whereas the in-plane vibrational modes only change by a factor of 3-4 over the same potential range. (We point out at this point that solution soluble isoquinoline would not exhibit this effect).

The explanation of the trends seen in figure 9 is that the out-of-plane vibrational modes have dipole derivative changes perpendicular to the metal surface when the molecules are lying flat on the surface. Absorption of radiation by the surface selection rule is thus allowed. When the orientation changes to the 6,7 configuration, absorption by these modes, which are now parallel to the surface is forbidden, as is any field induced interaction since most of the molecule lies outside of most of the field gradient. A large decrease in absorption (large increase in positive absorbance in the difference spectra) is thus expected and observed. In the case of the in-plane modes, the high electric field in the double layer leads to strong field induced absorption for the flatly adsorbed isoquinoline as expected. When reorienting to the 6,7 configuration, the modes become allowed by the surface selection rule. A smaller decrease in absorption is then observed.

CONCLUSIONS

Subtractively normalize interfacial Fourier transform infrared spectroscopy has been successfully used to follow the reorientations of isoquinoline molecules adsorbed onto a mercury electrode. It has been shown that field-induced infrared absorption makes a major contribution to the intensities of the vibrational band structure of aromatic organic molecules adsorbed on mercury.

The isoquinoline was observed to go through an abrupt reorientation at potentials more negative than about -0.73 V vs SCE (the actual transition potential being dependent on the bulk solution concentration) to the erect 6,7 standing position.

There was a lack of any potential dependence in the frequencies of the vibrational modes of the isoquinoline adsorbed to the mercury surface. The explanation of this may be the fact that mercury has no available vacant π - or δ -orbitals into which back-bonding (which is observed on all d metals) can occur. In this case, therefore, adsorption occurs only through σ -bonding. This point is under continued investigation.

ACKNOWLEDGEMENTS

We thank the Office of Naval Research for support of this work.

REFERENCES

1. A. Bewick and J. Robinson, J. Electroanal. Chem. 60 (1975) 163.
2. A. Bewick and J. Robinson, Surf. Sci. 55 (1976) 349.
3. R. Adzic, B.D. Cahan and E. Yeager, J. Chem. Phys. 58 (1973) 1780.
4. S. Pons, Ph.D. thesis, University of Southampton, England, 1979.
5. A. Bewick, J. M. Mellor and S. Pons, Electrochim. Acta. 25 (1980) 931.
6. W. G. Golden, D. S. Dunn and J. Overend, J. Catal. 71 (1981) 395.
7. S. Pons, J. Electroanal. Chem. 150 (1983) 495.
8. S. Pons and A. Bewick, Langmuir 1 (1985) 141.
9. C. Buess-Herman,, N. Vanlaethem-Meuree, G. Quarin, and L. Gierst, J. Electroanal. Chem. 123 (1981) 21.
10. M. W. Humphreys and R. J. Parsons, Electroanal. Chem. 82 (1977) 369.
11. R. Greef, in ref. 2.
12. Quarin, G., C. Buess-Herman, and L. Gierst, J. Electroanal. Chem. 123 (1981) 35.
13. C. Buess-Herman, J. Electroanal. Chem. 186 (1985) 27.
14. C. Buess-Herman, J. Electroanal. Chem. 186 (1985) 41.
15. C. Buess-Herman, C. Franck, C. and L. Gierst, Electrochim. Acta. 31 (1986) 965.
16. M.F. Crawford and R.E. MacDonald, Can. J. Phys. 36 (1958) 1022.
17. C. Korzeniewski, R.B. Shirts and S.Pons, J. Phys. Chem. 89 (1985) 2297.
18. C. Korzeniewski and S. Pons, Langmuir 2 (1986) 468.
19. C. Korzeniewski, University of Utah Ph.D Thesis (1987).
20. J.K. Foley, C. Korzeniewski, J.L. Daschbach, and S. Pons, "Electroanalytical Chemistry." Ed. A.J. Bard. Vol. 14. 309.
21. A. Bewick, private communication.

22. S.C. Wait, Jr. and J.C. McNerney, J. Mol. Spectrosc. 34 (1970) 56.

Table 1. A comparison of the characteristics of infrared vibrational bands arising from bulk solution species and adsorbed species.

BULK SOLUTION SPECIES	ADSORBED SPECIES
i) Bands appear with both s & p polarized light.	i) Bands appear with only p polarized light.
ii) Band positions independent of potential.	ii) Band positions may shift with potential.
iii) Relative intensity between bands is independent of potential.	iii) Relative intensity between bands may change with potential.
	iv) Some normally ir active bands may not be observed.
	iv) Some normally ir inactive bands may be observed.

Table 2. Assignments of the major infrared bands for isoquinoline at the mercury /water interface.

Band	Assignment	C _s	D _{2h}	In- or out-of-plane
1628	ν_8	A'	B _{3g}	In
1589	ν_9	A'	A _g	In
1575	$\nu_{26} + \nu_{38}$	A' + A''	A _g + B _{3u}	In + Out
1500	ν_{11}	A'	B _{2u}	In
1462	ν_{12}	A'	A _g	In
1435	ν_{13}	A'	B _{3g}	In
1380	ν_{14}	A'	A _g	In
1376	ν_{15}	A'	B _{1u}	In
1273	ν_{17}	A'	B _{1u}	In
1257	ν_{18}	A'	B _{2u}	In
1215	$\nu_{36} + \nu_{42}$	A'' + A''	A _u + B _{1g}	Out + Out
1180	ν_{19}	A'	B _{3g}	In


FIGURE LEGENDS

- Figure 1. SNIFTIRS difference spectra of the b_{3u} ring bending mode of p-difluorobenzene at a platinum electrode as a function of modulation potential.
- Figure 2. Possible orientations for the adsorption of isoquinoline on mercury.
- Figure 3. Superficial excess as a function of potential for a mercury electrode in contact with $0.5 \text{ mol dm}^{-3} \text{ Na}_2\text{SO}_4$ and the following isoquinoline concentrations: (a) 2.1×10^{-2} (saturated), (b) 6.3×10^{-3} , and (c) $2.1 \times 10^{-4} \text{ mol-dm}^{-3}$. (Data from Gierst et al., (1))
- Figure 4. Construction of the thin layer mercury reflectance cell used for all experiments.
- Figure 5. Schematic representation of the instrumentation used for SNIFTIRS experiments.
- Figure 6. Cyclic voltammogram for a mercury electrode in contact with a solution $2.1 \times 10^{-2} \text{ mol-dm}^{-3}$ in isoquinoline / $0.5 \text{ mol-dm}^{-3} \text{ Na}_2\text{SO}_4$ at 10 mV-s^{-1} .
- Figure 7. SNIFTIRS difference spectra for a mercury electrode in contact with a solution that is $1.3 \times 10^{-2} \text{ mol-dm}^{-3}$ in isoquinoline. Reference potential at 0.0 V vs SCE , sample potential (a) -0.60V , (b) -0.75V , (c) -0.80V and (d) -0.9V vs SCE .
- Figure 8. Expanded section of Figure 5 showing examples of differences in the magnitude of integrated area for two absorption bands with the same changes in electrode potential. The potentials for the difference spectra are -0.6 and -0.9V . The change in area for the 1380cm^{-1} in-plane mode has increased by a factor of ≈ 4 , whereas that of the 1215cm^{-1} out-of-plane mode has increased by a factor of ≈ 10 .


Figure 9. Plot of the normalized intensities of the bands observed in the SNIFTIRS difference spectra at a mercury electrode in contact with a $1.3 \times 10^{-2} \text{ mol-dm}^{-3}$ solution of isoquinoline vs the sample potential. Reference potential = 0.0V vs SCE.




Flat




4,5





6,7




5,6










$$\frac{\Delta R}{R}$$




WAVENUMBERS / cm^{-1}

20 cm^{-1}

SYMMETRY

$$\left[\frac{-\Delta R}{R} \right] (E=x)$$

$$\left[\frac{-\Delta R}{R} \right] (E=-0.1)$$



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. Robert A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, NY 14214

Dr. Douglas N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, UT 84602

Dr. Stanley Pons
~~Department of Chemistry~~
~~University of Utah~~
Salt Lake City, UT 84112

Dr. H. V. Venkatesetty
Honeywell, Inc.
10701 Lyndale Avenue South
Bloomington, MN 55420

Dr. J. Foos
EIC Labs Inc.
111 Downey St.
Norwood, MA 02062

Dr. Neill Weber
Ceramtec, Inc.
163 West 1700 South
Salt Lake City, UT 84115

Dr. Subhash C. Narang
SRI International
333 Ravenswood Ave.
Menlo Park, CA 94025

Dr. J. Paul Pemsler
Castle Technology Corporation
52 Dragon Ct.
Woburn, MA 01801

Dr. R. David Rauh
EIC Laboratory Inc.
111 Downey Street
Norwood, MA 02062

Dr. Joseph S. Foos
EIC Laboratories, Inc.
111 Downey Street
Norwood, Massachusetts 02062

Dr. Donald M. Schleich
Department of Chemistry
Polytechnic Institute of New York
333 Jay Street
Brooklyn, New York 01

Dr. Stan Szpak
Code 633
Naval Ocean Systems Center
San Diego, CA 92152-5000

Dr. George Blomgren
Battery Products Division
Union Carbide Corporation
25225 Detroit Rd.
Westlake, OH 44145

Dr. Ernest Yeager
Case Center for Electrochemical
Science
Case Western Reserve University
Cleveland, OH 44106

Dr. Mel Miles
Code 3852
Naval Weapons Center
China Lake, CA 93555

Dr. Ashok V. Joshi
Ceramtec, Inc.
2425 South 900 West
Salt Lake City, Utah 84119

Dr. W. Anderson
Department of Electrical &
Computer Engineering
SUNY - Buffalo
Amherst, Massachusetts 14260

Dr. M. L. Gopikanth
Chemtech Systems, Inc.
P.O. Box 1067
Burlington, MA 01803

Dr. H. F. Gibbard
Power Conversion, Inc.
495 Boulevard
Elmwood Park, New Jersey 07407

DL/1113/87/2

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. V. R. Koch
Covalent Associates
52 Dragon Court
Woburn, MA 01801

Dr. Randall B. Olsen
Chronos Research Laboratories, Inc.
4186 Sorrento Valley Blvd.
Suite H
San Diego, CA 92121

Dr. Alan Hooper
Applied Electrochemistry Centre
Harwell Laboratory
Oxfordshire, OX11 0RA UK

Dr. John S. Wilkes
Department of the Air Force
The Frank J. Seiler Research Lab.
United States Air Force Academy
Colorado Springs, CO 80840-6528

Dr. Gary Bullard
Pinnacle Research Institute, Inc.
10432 N. Tantan Avenue
Cupertino, CA 95014

Dr. J. O'M. Bockris
Ementech, Inc.
Route 5, Box 946
College Station, TX 77840

Dr. Michael Binder
Electrochemical Research Branch
Power Sources Division
U.S. Army Laboratory Command
Fort Monmouth, New Jersey 07703-5000

Professor Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton, Hants, SO9 5NH UK